No.



1900035

THE UNITED SHAMES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Speight Seed Farms, Inc.

Colhereus, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, therefore, this certificate of plant variety protection is to grant unto the said applicant(s) and the successors, heirs or assigns of the said applicant(s) for the term of seventeen years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic seed of the variety in a public repository as provided by LAW, the right to expect others from selling the variety, or offering it for sale, or reproducing it, orting it, or exporting it, or using it in producing a hybrid or different therefrom, to the extent provided by the Plant Variety Protection Act.

If therefrom, to the extent provided by the Plant Variety Protection Act.

A Certified seed and (2) shall conform to the number of generations

E OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

TOBACCO

'Speight G-70'

In Lestimony Winercot, I have hereunto set my hand and caused the seal of the Plant Variety Exotection Office to be affixed at the City of Washington

this 14th day of June in Se year of our Lord one thousand nine Lored and seventy-nine

Allast

Commissioner Plant Variety Protection Office

Agricultural Marketing Service

of Agriculture

| UNITED STATES DEPARTME AGRICULTURAL MARK | CETING SERVICE | | | FORM APPROVED OMB NO. 40-R3822 |
|---|--|--|--|---|
| LIVESTOCK, POULTRY, GRA APPLICATION FOR PLANT VARIE INSTRUCTIONS: See Reverse. | | | No certificate for pla be issued unless a co has been received (5) | ant variety protection may |
| 1a. TEMPORARY DESIGNATION OF VARIETY | 1b. VARIETY NAM | E | | AL USE ONLY |
| Speight G-70 | Speight | C-70 | 79000 | 25 |
| 2. KIND NAME | 3. GENUS AND SPE | | FILING DATE | TIME (A.M.) |
| Flue Cured Tobacco | Nicotiana | | 1-3-19 | TIME OLOO A.M. P.M. |
| 4. FAMILY NAME (BOTANICAL) Solanaceal Night Shade | 5. date of deter | 75 7810 | \$ 500.00 \$ A.50. 00 | 1-3-79 5-16-79 |
| 6. NAME OF APPLICANT(S) | 7. ADDRESS (Stree Code) | t and No. or R.F.D. No., | City, State, and ZIP | 8. TELEPHONE AREA |
| Speight Seed Farms, In | c. RFD 1 Winter | Box 507 ville, NC 2 | 8590 | 919 756-0718 |
| 9. IF THE NAMED APPLICANT IS NOT A PEORGANIZATION: (Corporation, partnersh Corporation | RSON, FORM OF ip, association, etc.) | 10. IF INCORPORAT DATE OF INCOR North Ca | ed, give state and Poration rolina | 11. DATE OF INCORPORATION 1-1-1972 |
| 12. NAME AND MAILING ADDRESS OF APPI | LICANT REPRESENTA | ATIVE(S), IF ANY, TO | SERVE IN THIS APPLIC | ATION AND RECEIVE |
| ALL PAPERS: M. L. Grimsle Speight Seed | • | | | |
| Winterville. | NC 28590 | | e e | |
| 13. CHECK BOX BELOW FOR EACH ATTACH | IMENT SUBMITTED: | | | |
| 13A. Exhibit A, Origin and Bree | ding History of the | Variety (See Section S | 52 of the Plant Variety | Protection Act.) |
| 🗵 13B. Exhibit B, Novelty Statem | ent. | • | | |
| 13C. Exhibit C, Objective Descr | iption of the Variety | (Request form from | Plant Variety Protecti | ion Office.} |
| X 13D. Exhibit D, Additional Desc | | | | · |
| 14a. DOES THE APPLICANT(S) SPECIFY THAT SEED? (See Section 83(a). (If "Yes," answer | T SEED OF THIS VAR er 14B and 14C below.) | ETY BE SOLD BY VAI | RIETY NAME ONLY AS | A CLASS OF CERTIFIED |
| 14b. DOES THE APPLICANT(S) SPECIFY THAT LIMITED AS TO NUMBER OF GENERATI | THIS VARIETY BE | 14c. IF "YES," TO 14 | B, HOW MANY GENERA | ATIONS OF PRODUC- |
| YES NO | | FOUNDATION | REGISTERED | X CERTIFIED |
| 15a. DID THE APPLICANT(S) FILE FOR PROTiname of countries and dates.) | ECTION OF THIS VAI | RIETY IN OTHER COU | NTRIES? TYES | NO (If "Yes," give |
| name of countries and dates.) | | | | |
| | | | e a tradition | |
| 15b. HAVE RIGHTS BEEN GRANTED THIS VA and dates.) | RIETY IN OTHER CO | UNTRIES? YES | NO (If "Yes," | give name of countries |
| | | 4 | | |
| | | | • | |
| 16. DOES THE APPLICANT(S) AGREE TO THE JOURNAL? YES | NO | | E(S) AND ADDRESS IN | |
| The applicant(s) declare(s) that a viable replenished upon request in accordance | sample of basic seed | d of this variety will b | e furnished with the a | pplication and will be |
| The undersigned applicant(s) is (are) the variety is distinct, uniform, and stable a 42 of the Plant Variety Act. | e owner(s) of this se | xually reproduced no | vel plant variety, and l | believe(s) that the provisions of Section |
| Applicant(s) is (are) informed that false | representation here | in can jeopardize pro | tection and result in pe | enalties. |
| 12-20-79 | | mark | 1 91 | P. Russ |
| (DATE) | | " IN COME (S | GIGNATURE OF APPLA | CANT) |
| 12 20 20 | ti da _n meter be | 011 | 1 21 | l = 1 |
| (DATE) | value sta | Kachel | SPECTE A | CANTO |

FORM GR-470 (1-78)

INSTRUCTIONS

GENERAL: Send an original copy of the application and exhibits, at least 2,500 viable seeds, and \$500 fee (\$250 filing fee and \$250 examination fee) to U.S. Dept. of Agriculture, Agricultural Marketing Service, Livestock, Poultry, Grain and Seed Division, Plant Variety Protection Office, National Agricultural Library Building, Beltsville, Maryland 20705. (See section 180.175 of the Regulations and Rules of Practice.) Retain one copy for your files. All items on the face of the form are self-explanatory unless noted below.

ITEM

- Give the date the applicant determined that he had a new variety based on (1) the definition in section 41(a) of the Act and (2) the date a decision was made to increase the seed.
- Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method; (2) the details of subsequent stages of selection and multiplication; (3) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified and (4) evidence of uniformity and stability.
- Give a summary statement of the variety's novelty. Clearly state how this novel variety may be distinguished from all other varieties in the same crop. If the new variety most closely resembles one or a group of related varieties:

 (1) identify these varieties and state all differences objectively; (2) attach statistical data for characters expressed numerically and demonstrate that these differences are significant; and (3) submit, if helpful, seed and plant specimens or photographs of seed and plant comparisons clearly indicating novelty.
- 13c Fill in the Exhibit C, Objective Description form, for all characteristics for which you have adequate data.
- Describe any additional characteristics that are not described, or whose description cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the description of characteristics that are difficult to describe, such as, plant habit, plant color, disease resistance, etc.
- If "YES" is specified (seed of this variety be sold by variety name only as a class of certified seed) the applicant may NOT reverse his affirmative decision after the variety has either been sold and so labeled, his decision published, or the certificate has been issued. However, if the applicant specified "NO," he may change his choice. (See section 180.16 of the Regulations and Rules of Practice.)
- See section 42 of the Plant Variety Protection Act and section 180.7 of the Regulations and Rules of Practice.

SPEIGHT SEED FARMS, INC.

BREEDERS, REGISTERED AND CERTIFIED SEED

TOBACCO SEED - SOYBEANS - HYBRID CORN 7900035

BOX 507 PHONE (919) 756-0718

WINTERVILLE, N. C. 28590

13-A

December 18, 1978

Exhibit A


Origin And Breeding History Of The Variety

Name: SPEIGHT G-70

Pedigree: Coker 258 x Va 115 x Speight G-10

- 1. In July, 1964, a cross was made between Va 115 and Speight G-10. Results of this cross was named or numbered G-16. After five years of individual plant selections from our disease nurseries, G-16 was found to be very highly resistant to Black Shank.
- 2. In the summer of 1969, Coker 258 was crossed on G-16 (C258 x G-16). Results of this cross was named or numbered N-61. Individual plant selections were made from N-61 through the sixth generation. The name of number N-61, was then changed to the variety name of Speight G-70.
- 3. Two variants were found during this five year period. One a stringy and narrow type, the other a very broad tall type with large fan shaped leaves. The type selected as N-61 was most like NC 2326. No variants were found in N-61 after the fourth generation.
- 4. Speight G-70 became stable in the fifth generation. It was tested in the NC Official Variety Test, by Speight Seed Farms and by farmers before entering the Regional Evaluation Program, G-70 remained stable throughout these tests.





SPEIGHT SEED FARMS, INC.

BREEDERS, REGISTERED AND CERTIFIED SEED TOBACCO SEED - SOYBEANS - HYBRID CORN

7900035

13-B

December 18, 1978

Novelty Statement

Name: Speight G-70----Flue Cured Tobacco Pediree: C-258 x Va 115 x Speight G-10

Speight G-70 most closely resembles NC 2326, but differs by growing shorter (38.5 vs. 41.1 inches) in height. It has a higher leaf count (19.0 vs 18.4) leaves per plant; leaf sizes and shapes are about the same. Internodes average shorter (2.4 vs 2.6 inches); Ground suckers counts are higher (0.7 vs 0.2). G-70 flowers three or four days later (64 vs 60 days) and flower heads are more closed (intermediate vs open). G-70 is resistant to Root Knot Nemotode---NC 2326 is susceptible. G-70 normally yields about 500 pounds per acre more and standibility is better-due to bigger and stronger root system.

Information in 1977 and 1978 Flu-Cured Tobacco Variety Evaluation Report

Disease Data From Regional Small Plot Test 1977 and 1978

| Entry of No. | B.S. Index | Root Knot | B. Wilt Index | B. Spot Tolerance |
|--------------|---------------|--------------|------------------|----------------------|
| NC 95 | 31.2 | Res. | 29.3 | Tolerant |
| NC 2326 | 33.5 | Sus. | 53.0 | Mod. Tolerant |
| Speight G-70 | 19.0 | Res. | 40.1 | Mod. Tolerant |






Table 1. (Continued)
Whiteville, Kinston, Oxford, Blackstone, Tifton and Florence - 1977

| Entry | Sol. Sug. % | Tot. Nit. % | Ins. Nit. % | Alpha Amino N. % | Plant Height Inches | Leaf Number | Days to Flower |
|---------|-------------------|-------------------|-------------------|------------------------|---------------------------|----------------|----------------------|
| NC 2326 | 11.3 | 2.67 | 1.08 | .32 | 41.1 | 18.4 | 60 |
| NC 95 | 11.1 | 2.76 | 1.14 | .32 | 40.4 | 19.0 | 63 |
| 3 | 11.1 | 2.71 | 1.16 | .31 | 37.1 | 18.6 | 66 |
| 4 | 11.2 | 2.66 | .92- | .30 | 38.2 | 18.5 | 64 |
| 5 | 11.3 | 2,59 | 1.13 | .31 | 40.0 | 20.2 | 63 |
| 6 | 9.8 | 2.87 | 1.12 | .34 | 45.2 | 19.0 | 62 |
| 7 | 10.8 | 2.68 | 1.10 | .29 | 41.6 | 20.5 | 67 |
| 8 | 10.5 | 2.79 | 1.16 | .32 | 42.1 | 20.7 | 68 |
| 9 | 11.8 | 2.69 | 1.08 | .33 | 38.9 | 17.9 | 63 |
| 10 | 10.7 | 2.89 | 1.17 | .34 | 43/.1 | 18.4 | 63 |
| 11 | 11.9 | 2.51 | 1.06 | .26 | 40.9 | 21.7 | 71 |
| 12 | 11.1 | 2.68 | 1.14 | .31 | 43.7 | 18.9 | 61 |
| 13 | 11.9 | 2.59 | 1.09 | .30 | 43.6 | 18.9 | 65 |
| 14 | 11.0 | 2.74 | 1.16 | .31 | 45.5 | 20.2 | 65 |
| 15 | 11.5 | 2.77 | 1.13 | .32 | 39.0 | 18.8 | 59 |
| 16 | 11.2 | 2.74 | 1.17 | .34 | 42.1 | 19.9 | 64 |
| . 17 | 10.8 | 2.78 | 1.26+ | .31 | 42.1 | 22.1 | 70 |

TURN TO NEXT PAGE- Plot 30 = Speight G-70

Table (Continued)
Whiteville, Kinston, Oxford, Blackstone, Tifton and Florence - 1977

| Entry | Sol. Sug. % | Tot. Nit. % | Ins. Nit. % | Alpha Amino N. % | Plant Height Inches | Leaf Number | Days to Flower |
|-----------|-------------------|-------------------|-------------------|------------------------|---------------------------|----------------|----------------------|
| 18 | 11.0 | 2.70 | 1.18 | .32 | 42.8 | 22.4 | 71 |
| 19 | 11.4 | 2.64 | 1.07 | .27 | 42.0 | 20.2 | 63 |
| 20 | 10.9 | 2.60 | 1.13 | .30 | 39.9 | 20.9 | 65 |
| 21 | 10.8 | 2.69 | 1.13 | .32 | 41.0 | 19.8 | 62 |
| 22 | 10.8 | 2.82 | 1.14 | .35 | 39.9 | 19.0 | 61 |
| 23 | 10.5 | 2.60 | 1.17 | .35 | 41.1 | 21.9 | 67 |
| 24 | 10.8 | 2.71 | 1.14 | .30 | 41.6 | 20.9 | 66 |
| 25 | 10.5 | 2.92 | 1.25+ | .31 | 45.1 | 19.6 | 65 |
| 26 | 11.4 | 2.54 | 1.09 | .31 | 42.3 | 19.6 | 65 |
| 27 | 9.8 | 2.88 | 1.18 | .30 | 42.2 | 21.7 | 70 |
| 28 | 9.8 | 2.88 | 1.23+ | .39+ | 40.5 | 18.7 | 60 |
| 29 | 10.5 | 2.66 | 1.14 | ,32 | 36.2 | 20.3 | 61 |
| 30 G - 20 | 11.7 | 2.60 | 1.12 | .31 | 38.5 | 19.0 | 64 |
| 31 | 11.6 | 2.40- | 1.04 | .28 | 44.9 | 22.6 | 68 |
| 32 | 9.5- | 2.91 | 1.23+ | .35 | 40.3 | 20.5 | 63 |
| 33 | 11.6 | 2.67 | 1.17 | .32 | 44.0 | 19.6 | 67 |

13-e bil-e

UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE

OBJECTIVE DESCRIPTION OF VARIETY

Tobacco (Nicotiana tabacum) VARIETY NAME OR TEMPORARY DESIGNATION NAME OF APPLICANT(S) Speight Seed Farms, Inc. 6-70 Speight ADDRESS (Street and No. or R.F.D. No., City, State, and Zip Code) FOR OFFICIAL USE ONLY PVPO NUMBER RFD 1 Box 507 7900035 Winterville, NC 28590 Place the appropriate number that describes the varietal character in the boxes below. [0]9] when number is either 99 or less or 9 or less. Place a zero in first box (e.g. 0 8 9 or 1. CLASS: 6 = CIGAR WRAPPER 5 = CIGAR BINDER 4 = CIGAR FILLER 3 = AIR-CURED 2 = FIRE-CURED 1 = FLUE-CURED 9 = FOREIGN-NON-CIGAR LEAF 8 = FOREIGN-CIGAR LEAF 7 = MISCELLANEOUS-DOMESTIC 3 = DARK AIR-CURED 2 = MARYLAND AIR-CURED: 1 = BURLEY STANDARD VARIETIES 06 = SC 5803 = COKER 319 04 = HICKS 05 = SPEIGHT G-280 2 = NC 2326 01 = NC 95 12 = Ky 165 10 = Ky 10 11 = MARYLAND 609 08 = BURLEY 21 09 = BURLEY 49 07 = Ky 15116 = OTHER 15 = FLORIDA 17 13 = Pennoel 69 14 = HAVANA 503 2. MATURITY (Transplant to 50% plants 1 Fl.) (Select code from Standard Varieties listed above) DAYS EARLIER THAN . NO, OF DAYS 6 DAYS LATER THAN . . 0 3. SEEDING TO TRANSPLANTING (Select code from Standard Varieties listed above) DAYS EARLIER THAN 8 NO. OF DAYS DAYS LATER THAN ... 0 4. PLANT HEIGHT (After topping) (Select code from Standard Varieties listed above) CM SHORTER THAN. 0 9 CM TALL CM TALLER THAN . 5. LEAF SIZE (At leaf maturity) (Select code from Standard Varieties listed above) LENGTH CM 15TH LEAF CM 10TH LEAF CM 5TH LEAF 6 6 8 5 8 CM SHORTER THAN CM SHORTER THAN; 9 CM SHORTER THAN . 2 0 0 CM LONGER THAN . 0 2 2 2 2 CM LONGER THAN . 0 CM LONGER THAN . 8 WIDTH CM 15TH LEAF CM 5TH LEAF 2 7 CM 10TH LEAF 3 8 1 CM NARROWER THAN CM NARROWER THAN. CM NARROWER THAN . CM WIDER THAN 2 CM WIDER THAN 0 2 CM WIDER THAN . . . 6. LEAF YIELD (Select code from Standard Varieties listed above) % MORE THAN ... % LESS THAN ... 3

| | | | | CTANDARD \ | ADIETIES | | 700011357 7 |
|--------------------------|------------------------|--------------------------|-----------------------------|-----------------------|-------------------------------|----------------------|---------------------------------------|
| GROUPIN | 7 | = NC 2326 | 03 = COKER 319 | STANDARD \ | | IGHT G-28 | 7900035 06 = SC 58 |
| 01 = NC 95 07 = Ky 15 | | = NC 2320 = BURLEY 21 | 09 = BURLEY 49 | 10 = Ky 10 | - ·· | RYLAND 609 | 12 = Ky 165 |
| 13 = Pennb | | = HAVANA 503 | 15 = FLORIDA 17 | 16 = OTHE | | | |
| | | | andard Varieties listed abo | ········· | | | |
| | D NORMAL | | andard Addiestes tisted and | 140) | | | |
| 1 9 | | ER PLANT | | | | | |
| 1 9 NOT TO | O NO. O | F LEAVES > 4 | 0.6 CM | | 9 8 см н | EIGHT OF LAS | T LEAF> 40.6 CM |
| 2 4 | 0 NO. O | F LEAVES OR NO | DES TO "CROWFOOD" | FROM 1ST H | ARVESTABLE LE | AF | |
| 8. INTER | NODES (Top | ped) (Select code | from Standard Varieties li | sted above) | | | · · · · · · · · · · · · · · · · · · · |
| 6 1 | MM LENG | этн (|) 5 MM SHORTER | THAN | 0 2 | MM LO | NGER THAN |
| and the second | CHARACTEI LE ANGLE: | | | | | | |
| 3 5 | DEGREE | s 1 | GROUPING: 1 = < | 35 ⁰ 2 = | 35-45 ⁰ 3 = | = 46–65 ⁰ | 4 = > 65° |
| LEAF | CARRIAGE | No. of the second | | LEA | F COLOR (At top | ping or when 50 | % of plants with 1 flower) |
| 2 | 3 = UPRIGH |) (DROOPING) T | 2 = HORIZONTAL | <u>_</u> 3 | 1 = LIGHT GRE 4 = YELLOW-C | | EEN 3 = DARK GREEN = YELLOW |
| LEAF : | SHAPE: | | | | | | |
| 2 | Park Contract | R THAN LONG | 2 = LENGTH EQUALS | width $oxedsymbol{1}$ | 1 = BROADEST 3 = ABOVE MII | | 2 = BELOW MIDDLE |
| TIPSH | IAPE | | | VEN | IATION PATTER | N: | |
| 2 | 1 = ACUTE | 2 = ACUMI | NATE 3 = OBTUSE | 2 | 1 = SQUARE | 2 = ANO | SULAR |
| LEAF S | SURFACE | | | LEA | F MARGIN | 1 | |
| 1 | 1 = SMOOTH | ı (HICKS) | 2 = PUCKERED (NO | 95) 2 | 1 = WAVY 2 | = NOT WAVY | 1 = RECURVED 2 = NOT RECURVED |
| 10. FLOW | /ERS: | | | FLC | WER HEAD HAB | IT: | |
| 2 | COLOR 1= | | = PINK | 2 | 1 = CLOSED (N | Annual Control | 2 = INTERMEDIATE |
| 44 DI 482 | 3 = T CODM | RED 4 | = OTHER | L | 3 = OPEN (HIC | KS) | |
| 11. PLAN | TFORM | | | | | | |
| 2 1 | 1 = PYRAMII | DAL 2=0 | COLUMNAR 3= | OTHER (Spec | ify) | | |
| 12. GROU | IND SUCKE | RS: | | | | | |
| 0.7 | NO, PER PLA | ANT | | | | | |
| 13. DISEA | ASE (O = Not | t tested, 1 = Suscep | tible, 2 = Resistant) | | | | |
| 2 | BLACK SHA | NK (RACES) | | [2] | FUSARIUM WI | LT (NICOTIAN | A) |
| 2 | BLACK ROO | твот | | 0 | FUSARIUM WI | LT (BATATAS) | |
| 0 • | BLUE MOLD | | | 0 | FROGEYE | | |
| 0 | WILDFIRE (5 | SPECIES) | | 2 | BROWN SPOT | Tolerance | |
| | BLACKFIRE | | | 2 | BACTERIAL W | ILT | |

NC

2326

3

NC 2326

| CHARACTER | VARIETY | D FOR THE CHARACTERS GIVEN: | VARIETY |
|--------------|-----------|-----------------------------|---------|
| MATURITY | Coker 319 | LEAF TIP SHAPE | NC 2326 |
| EAF LENGTH | NC 2326 | VENATION PATTERN | NC 2326 |
| EAF WIDTH | NC 2326 | LEAF SURFACE | NC 2326 |
| EAF CARRIAGE | NC 2326 | LEAF MARGIN | NC 2326 |
| ETIOLE ANGLE | NC 95 | LEAF COLOR | NC 2326 |
| LEAF SHAPE | NC 2326 | PLANT FORM | NC 95 |

16. COMMENTS (For increasing accuracy of description)

2326

5

NC

SUBMITTED

STANDARD

NAME OF STANDARD

VARIETY

Speight G-70 most closely resembles NC 2326 but differs by growing shorter (38.5 vs 41.1 inches) with higher leaf counts (19.0 vs 18.4) leaves per plant. Leaf size and shapes are abouth the same as NC 2326. Internodes average shorter (2.4 vs 2.6 inches). General sucker counts are higher (0.7 vs 0.2). G-70 flowers about three or four days later (64 vs 60 days) from transplanted date. G-70 carries Root Knot nemotode resistance and NC is susceptible. Flower heads are more closed; intermediate, for G-70 and open for NC 2326.

Disease Data From Reg. Small Plot Test And NC Off. Var. Test 1978

| | B.S. | Bac. Wilt | R. Kn. | Brown Spot |
|---------|-------|-----------|--------|---------------|
| Entry | Index | Index | Nem. | Tolerance |
| NC 2326 | 33.5 | 53.0 | Sus. | Mod. Tolerant |
| NC 95 | 31.2 | 29.3 | Res. | Tolerant |
| G- 70 | 19.0 | 40.1 | Res. | Mod. Tolerant |

2326

nc

SPEIGHT SEED FARMS, INC.

BREEDERS, REGISTERED AND CERTIFIED SEED
TOBACCO SEED - SOYBEANS - HYBRID CORN

BOX 507 ● PHONE (919)756-0718 WINTERVILLE, N. C. 28590

13-D

December 18, 1978

Exhibit D

Speight G-70 Flue Cured Tobacco

Speight G-70 flowers four days later than NC 2326 but is similar in many ways. G-70 starts off growing flatter and wider, it closes in the middle faster. NC 2326 seems to grow more erect and fills the middle later. Both grow with dark green color. G-70 normally produces about one more leaf per plant. It grows a larger and stronger root system; standibility and storm resistance are better. The topped out height of G-70 is usually one or two inches shorter. Tops differ by being more closed on G-70, intermediate vs open, with three inches more distance between crows foot and top leaf on NC 2326. Leaves are closer spaced especially in the top part of the plants. They are smooth and hang parallel on both varieties. Expect a few more ground suckers on G-70 with less leaf axil suckers. G-70 normally yields 500 pounds per acre more than NC 2326. Color of cured leaf is about the same with, G-70 producing a higher percentage of medium to heavy bodied, and less chaffy leaf. Nicotine content is usually lower. G-70 carries very high resistance to Root Knot Nemotode. NC 2326 carries moderate resistance to Black Shank, low to Bacterial Wilt and no resistance to Root Knot Nemotodee

1978 Comparison in Regional Small Plot & Regional Farm Test

| Entry or No. | Yield lbs/A | Value \$/A | Index \$ CWT | Nic.% | R. Kn. Res. |
|--------------|-------------|------------|--------------|-------|-------------|
| NC 2326 | 2808 | \$3840 | \$136.52 | 3.09 | Sus. |
| NC 95 | 2684 | \$3524 | \$131.36 | 3.32 | Res. |
| Speight G-70 | 3256 | \$4414 | \$134.98 | 2.76 | Res. |

Inf. in Flue Cured Tobacco Evaluation Committee Report 1977-1978





9

SPEIGHT G-70